

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM	
1. REPORT NUMBER		RECIPIENT'S CATALOG NUMBER	
AFOSR/TR-80-1164		AD A092191	
4. TITLE (and Subtitle)		5. TYPE OF REPORT & PERIOD COVERED	
ASYMPTOTIC DISTRIBUTIONS OF FUNCTIONS OF THE EIGENVALUES OF THE REAL AND COMPLEX NONCENTRAL WISHART MATRICES.		Interim rept.	
7. AUTHOR(s)		6. PERFORMING ORG. REPORT NUMBER	
C. Fang and P. R. Krishnaiah		F49620-79-C-0161	
9. PERFORMING ORGANIZATION NAME AND ADDRESS		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS	
University of Pittsburgh Department of Mathematics and Statistics Pittsburgh, PA 15260		61102F 2304/AS	
11. CONTROLLING OFFICE NAME AND ADDRESS		12. REPORT DATE	
Air Force Office of Scientific Research /NM Bolling AFB, Washington, DC 20332		11 July 1980	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES	
15 TR-80-11		39	
16. DISTRIBUTION STATEMENT (of this Report)		15. SECURITY CLASS. (of this report)	
Approved for public release; distribution unlimited		UNCLASSIFIED	
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
		E	
18. SUPPLEMENTARY NOTES			
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)			
Asymptotic Distributions, Noncentral Wishart Matrix, Functions of Eigenvalues Complex Matrices, Mixtures of Populations, Quadratic Forms, Perturbation Theory			
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)			
In this paper, the authors derived asymptotic expressions for the joint densities of various functions of the noncentral real and complex Wishart matrices. Applications of the above results are also discussed in the area of reduction of dimensionality and testing for the structure of interaction in two-way classification with one observation per cell.			

ASYMPTOTIC DISTRIBUTIONS OF FUNCTIONS  
OF THE EIGENVALUES OF THE REAL AND  
COMPLEX NONCENTRAL WISHART MATRICES\*

C. Fang  
University of Pittsburgh  
and  
Carnegie-Mellon University

P. R. Krishnaiah  
University of Pittsburgh

Accession For	
NIIS GNA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	

July 1980

Technical Report No. 80-11

Institute for Statistics and Applications  
Department of Mathematics and Statistics  
University of Pittsburgh  
Pittsburgh, PA. 15260

\*This work is sponsored by the Air Force Office of Scientific Research, Air Force Systems Command under Contract F49620-79-C-0161. Reproduction in whole or in part is permitted for any purpose of the United States Government.

80 11 06 033

Approved for public release;  
distribution unlimited.

## 1. Introduction

The distributions of functions of the eigenvalues of the real and complex Wishart matrices are very useful in studying the structures of the covariance matrices of the real and complex multivariate normal distributions respectively and other problems. Krishnaiah and Lee (1977) derived the joint asymptotic distributions of the linear functions as well as the ratios of the linear functions of the roots of the central Wishart matrix when the population covariance matrix has simple roots. Fujikoshi (1978) derived an asymptotic expression for the distribution of a function of the roots of the central Wishart matrix when the roots have multiplicity whereas Krishnaiah and Lee (1979) obtained corresponding expressions for the joint density of the functions of the roots. In this paper, we obtain asymptotic expressions for the joint densities of various functions of the noncentral real and complex Wishart matrices. These expressions are in terms of multivariate normal density and multivariate Hermite polynomials. Percentage points of some test statistics are computed by using the above asymptotic expressions and these percentage points are compared with the results obtained by simulation. Applications of the above results are also discussed in problems of studying the structure of interactions, mixtures of multivariate normal populations, and reduction of dimensionality. The results obtained on the joint distribution of the functions of the eigenvalues of the real Wishart matrix are generalized to the case of multivariate quadratic forms. Finally, the joint asymptotic distribution of the functions of the roots of the complex Wishart matrix is derived.

## 2. Perturbation Technique

Let  $\ell_1 \geq \dots \geq \ell_p$  be the eigenvalues of the symmetric matrix  $T: p \times p$ , and  $\lambda_1 \geq \dots \geq \lambda_p$  are the eigenvalues of the symmetric matrix  $V: p \times p$ , where

$$T(\epsilon) = V + \epsilon V^{(1)} + \epsilon^2 V^{(2)} + \dots \quad (2.1)$$

Then, there exist orthogonal matrices  $\Gamma$  and  $G$  such that  $T = GLG'$  and  $V = \Gamma \Lambda \Gamma'$ , where  $L = \text{diag}(\ell_1, \dots, \ell_p)$ ,  $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_p)$ . The columns of  $\Gamma$  and  $G$  consist of the eigenvectors of  $V$  and  $T$  respectively.

Lawley (1956), Mallows (1961), Izenman (1972) and Fujikoshi (1978) have approximated the eigenvalues and eigenvectors of  $T$  in various papers. The authors have either assumed that  $\lambda_1$ 's do not have multiplicity or the approximations were established by tacitly assuming that the eigenvalues and eigenvectors admit series expansions in the infinitesimal parameter  $\epsilon$  as follows:

$$\ell_j = \lambda_j + \epsilon \lambda_j^{(1)} + \epsilon^2 \lambda_j^{(2)} + \dots \quad (2.2)$$

$$\tilde{g}_j = \Gamma_j + \epsilon \tilde{\Gamma}_j^{(1)} + \epsilon^2 \tilde{\Gamma}_j^{(2)} + \dots$$

and no attempts were made to prove that the series converge. A more insight treatment to settle this question of convergence is found in Kato (1976).

Now  $T(\epsilon)$  and  $V$  are linear transformations which operate on the  $p$ -dimensional complex vector field  $C^p$ ,  $\epsilon$  is also complex,  $\lambda_1 \geq \dots \geq \lambda_p$  are the eigenvalues of  $V: p \times p$  such that

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)  
2 NOTICE OF TRANSMITTAL TO DDC  
This technical report has been reviewed and is  
approved for public release IAW AFR 190-12 (7b).  
Distribution is unlimited.  
A. D. BLOSE  
Technical Information Officer

$$\lambda_{q_1+\dots+q_{\alpha-1}+1} = \dots = \lambda_{q_1+\dots+q_{\alpha}} = \theta_{\alpha} \quad (2.3)$$

for  $\alpha = 1, 2, \dots, r$ ,  $q_1 + \dots + q_r = p$ ,  $q_0 = 0$  and let

$J_{\alpha}$  ( $\alpha=1, \dots, r$ ) denote the set of integers

$q_1 + \dots + q_{\alpha-1} + 1, \dots, q_1 + \dots + q_{\alpha}$ . We need the following lemma in the sequel.

Lemma 2.1. For Hermitian matrices  $T(\epsilon)$  and  $V$  as defined before,

$$T(\epsilon) = V + \epsilon V^{(1)} + \epsilon^2 V^{(2)} + \dots$$

and  $V$  is diagonalized as  $V = \text{diag}(\lambda_1, \dots, \lambda_p)$ . Then the mean eigenvalue of  $T(\epsilon)$  corresponding to  $\theta_{\alpha}$  which is the eigenvalue of  $V$  with multiplicity  $q_{\alpha}$ , is

$$\bar{\lambda}_{\alpha}(\epsilon) = \theta_{\alpha} + \epsilon \bar{\lambda}_{\alpha}^{(1)} + \epsilon^2 \bar{\lambda}_{\alpha}^{(2)} + \dots \quad (2.4)$$

where

$$\begin{aligned} \bar{\lambda}_{\alpha}^{(1)} &= \frac{1}{q_{\alpha}} \text{tr} V_{\alpha\alpha}^{(1)} \\ \bar{\lambda}_{\alpha}^{(2)} &= \frac{1}{q_{\alpha}} \text{tr} [V_{\alpha\alpha}^{(2)} + \sum_{\beta \neq \alpha} (\theta_{\alpha} - \theta_{\beta})^{-1} V_{\alpha\beta}^{(1)} V_{\beta\alpha}^{(1)}] \\ \theta_{\alpha\beta} &= \theta_{\alpha} - \theta_{\beta} \end{aligned} \quad (2.5)$$

with

$$V^{(i)} = \begin{pmatrix} V_{11}^{(i)} & V_{12}^{(i)} & \dots & V_{1r}^{(i)} \\ \vdots & & & \\ \vdots & & & \\ V_{r1}^{(i)} & V_{r2}^{(i)} & \dots & V_{rr}^{(i)} \end{pmatrix}$$

and  $V_{\alpha\beta}^{(1)}$  is of order  $q_\alpha \times q_\beta$ .

When  $q_1 = \dots q_r = 1$ , the above lemma was proved in Lawley (1956). When  $q_\alpha \geq 1$  ( $\alpha=1, \dots, r$ ), the lemma was given implicitly in Kato (1976). For  $q_\alpha = 1$  the normalized eigenvector of  $T(\epsilon)$  corresponding to  $\theta_\alpha$  is  $G_\alpha(\epsilon) = (G_{1\alpha}(\epsilon), \dots, G_{p\alpha}(\epsilon))'$ , with

$$G_{i\alpha}(\epsilon) = \frac{a_{i\alpha}(\epsilon)}{(\theta_\alpha - \lambda_1)} + \sum_{j \neq \alpha} \frac{a_{ij}(\epsilon)a_{j\alpha}(\epsilon)}{(\theta_\alpha - \lambda_1)(\theta_\alpha - \lambda_j)} - \frac{a_{i\alpha}(\epsilon)a_{\alpha\alpha}(\epsilon)}{(\theta_\alpha - \lambda_1)^2} + \dots i \neq \alpha$$

$$G_{\alpha\alpha}(\epsilon) = 1 - \frac{1}{2} \sum_{j \neq \alpha} \frac{a_{i\alpha}(\epsilon)a_{\alpha i}(\epsilon)}{(\theta_\alpha - \lambda_j)(\theta_\alpha - \lambda_j)} + \dots \quad (2.6)$$

where

$$A(\epsilon) = T(\epsilon) - V = (a_{ij}(\epsilon))$$

The series (2.4), (2.6) are convergent for

$$|\epsilon| < \left( \frac{2c_1}{d} + c_2 \right)^{-1} \quad (2.7)$$

where  $c_1, c_2 \geq 0$  such that  $||V^{(j)}|| \leq c_1 c_2^{j-1}$  for  $j = 1, 2, \dots$ , and  $d = \min(|\theta_\alpha - \theta_{\alpha-1}|, |\theta_\alpha - \theta_{\alpha+1}|)$ .

### 3. Asymptotic Joint Distributions of Functions of the Roots of Noncentral Wishart Matrix

Let the columns of  $X: p \times n$  be distributed as multivariate normal with covariance matrix  $\Sigma = (\sigma_{ij})$  and means given by  $E(X) = U = (\mu_1, \dots, \mu_n)$ , where  $\mu'_j = (\mu_{j1}, \dots, \mu_{jp})$ . Then,  $S = XX' = (S_{ij})$  is distributed as the central or noncentral Wishart matrix  $W_p(n, \Sigma, M)$  with  $n$  degrees of freedom according as  $M = 0$  or  $M \neq 0$  where  $M = \sum_{j=1}^n \mu_j \mu'_j = n(v_{ij})$ . Now, let  $\lambda_1 \geq \dots \geq \lambda_p$  denote the eigenvalues of  $S/n$  whereas  $\lambda_1 \geq \dots \geq \lambda_p$  denote the eigenvalues of  $E(S/n) = \Sigma + M/n = \Lambda$ . Without loss of generality, we assume that  $\Lambda = \text{diag.} (\lambda_1, \dots, \lambda_p)$ . Also, let

$$\lambda_{q_1 + \dots + q_{\alpha-1} + 1} = \dots = \lambda_{q_1 + \dots + q_{\alpha}} = \theta_{\alpha} \quad (3.1)$$

for  $\alpha = 1, 2, \dots, r$ ,  $q_1 + \dots + q_r = p$ , and  $q_0 = 0$ .

In this section, we derive the joint asymptotic distribution of  $L_1, \dots, L_k$  where  $L_j = \sqrt{n} \{T_j(\ell_1, \dots, \ell_p) - T_j(\lambda_1, \dots, \lambda_p)\}$  and  $T_j(\ell_1, \dots, \ell_p)$  satisfy the following assumptions:

(i)  $T_j(\ell_1, \dots, \ell_p)$  is analytic about  $\lambda_1, \dots, \lambda_p$

$$(ii) \quad \left. \frac{\partial T_i(\ell_1, \dots, \ell_p)}{\partial \ell_{j_1}} \right|_{\ell=\lambda} = c_{ij_1} = a_{i\alpha}$$

$$\left. \frac{\partial^2 T_i(\ell_1, \dots, \ell_p)}{\partial \ell_{j_2} \partial \ell_{j_1}} \right|_{\ell=\lambda} = c_{ij_1 j_2} = a_{i\alpha\beta}$$

$$\left. \frac{\partial^3 T_i(\ell_1, \dots, \ell_p)}{\partial \ell_{j_3} \partial \ell_{j_2} \partial \ell_{j_1}} \right|_{\ell=\lambda} = c_{ij_1 j_2 j_3} = a_{i\alpha\beta\gamma} \quad (3.2)$$

for  $j_1 \in J_\alpha$ ,  $j_2 \in J_\beta$ ,  $j_3 \in J_\gamma$ ,  $\lambda' = (\lambda_1, \dots, \lambda_p)$ ,  $\ell' = (\ell_1, \dots, \ell_p)$   
and  $J_\alpha$  denotes the set of integers  $q_1 + \dots + q_{\alpha-1} + 1, \dots, q_1 + \dots + q_\alpha$   
for  $\alpha = 1, 2, \dots, r$ .

Expanding  $T_i(\ell_1, \dots, \ell_p)$  as the Taylor series, we obtain

$$\begin{aligned} T_i(\ell_1, \dots, \ell_p) &= T_i(\lambda_1, \dots, \lambda_p) + \sum_{\alpha=1}^r a_{i\alpha} \sum_{j_1 \in J_\alpha} (\ell_{j_1} - \theta_\alpha) \\ &+ \frac{1}{2} \sum_{\alpha=1}^r \sum_{\beta=1}^r a_{i\alpha\beta} \sum_{j_1 \in J_\alpha} \sum_{j_2 \in J_\beta} (\ell_{j_1} - \theta_\alpha)(\ell_{j_2} - \theta_\beta) \\ &+ \frac{1}{6} \sum_{\alpha\beta\gamma} a_{i\alpha\beta\gamma} \sum_{j_1 \in J_\alpha} \sum_{j_2 \in J_\beta} \sum_{j_3 \in J_\gamma} (\ell_{j_1} - \theta_\alpha)(\ell_{j_2} - \theta_\beta)(\ell_{j_3} - \theta_\gamma) \\ &+ \text{terms of higher degree.} \end{aligned} \quad (3.3)$$

Now, let

$$S/n = \Lambda + \frac{1}{\sqrt{n}} V \quad (3.4)$$

where

$$V = \sqrt{n} \left( \frac{S}{n} - \Lambda \right) = \begin{pmatrix} v_{11} & v_{12} & \dots & v_{1r} \\ v_{21} & v_{22} & \dots & v_{2r} \\ \vdots & \vdots & & \vdots \\ v_{r1} & v_{r2} & \dots & v_{rr} \end{pmatrix}$$

So by applying Lemma 2.1 on (3.4), we obtain

$$L_i = \sum_{\alpha=1}^r a_{i\alpha} \operatorname{tr} Z_{\alpha}^{(1)} + \frac{1}{\sqrt{n}} \left\{ \sum_{\alpha=1}^r a_{i\alpha} \operatorname{tr} Z_{\alpha}^{(2)} + \frac{1}{2} \sum_{\alpha=1}^r \sum_{\beta=1}^r a_{i\alpha\beta} \right. \\ \left. (\operatorname{tr} Z_{\alpha}^{(1)}) (\operatorname{tr} Z_{\beta}^{(1)}) \right\} + O(n^{-1}) \quad (3.5)$$

where  $Z_{\alpha}^{(1)} = V_{\alpha\alpha}$ ,  $Z_{\alpha}^{(2)} = \sum_{\beta \neq \alpha} \theta_{\alpha\beta}^{-1} V_{\alpha\beta} V_{\beta\alpha}$  and  $\theta_{\alpha\beta} = \theta_{\alpha} - \theta_{\beta}$ .

Also,

$$\operatorname{tr} Z_{\alpha}^{(1)} = \sqrt{n} \sum_{j_1 \in J_{\alpha}} \left( \frac{S_{j_1 j_1}}{n} - \lambda_{j_1} \right), \\ \operatorname{tr} Z_{\alpha}^{(2)} = \frac{1}{n} \sum_{\beta \neq \alpha} \theta_{\alpha\beta}^{-1} \sum_{j_1 \in J_{\alpha}} \sum_{j_2 \in J_{\beta}} S_{j_1 j_2}^2.$$

After some algebraic manipulations, we obtain the following expression for the joint characteristic function of  $L_1, \dots, L_k$ :

$$\begin{aligned} \psi(t_1, \dots, t_k) &= E\left\{\exp\left(i \sum_{j=1}^k t_j L_j\right)\right\} \\ &= E\left[\exp\left(i \sum_{j=1}^k \sum_{\alpha=1}^r t_j a_{j\alpha} \operatorname{tr} Z_{\alpha}^{(1)}\right) \right. \\ &\quad \times \left\{1 + \frac{1}{\sqrt{n}} \left(i \sum_{j=1}^k \sum_{\alpha=1}^r t_j a_{j\alpha} \operatorname{tr} Z_{\alpha}^{(2)} + \frac{i}{2} \sum_{j=1}^k \sum_{\alpha=1}^r \sum_{\beta=1}^r t_j a_{j\alpha\beta} \right. \right. \\ &\quad \left. \left. \operatorname{tr} Z_{\alpha}^{(1)} \operatorname{tr} Z_{\beta}^{(1)}\right) \right\} \left. \right] = E_1(\underline{t}) + E_2(\underline{t}) + E_3(\underline{t}) + O(n^{-1}) \end{aligned} \quad (3.6)$$

where  $\underline{t}' = (t_1, \dots, t_k)$ . In Eq. (3.6),

$$E_1(\underline{t}) = E\left[\exp\left(i \sum_{i=1}^k \sum_{\alpha=1}^r t_i a_{i\alpha} \operatorname{tr} Z_{\alpha}^{(1)}\right)\right]$$

$$= \text{etr}(-i \sqrt{n} B) |I - 2i B \Sigma / \sqrt{n}|^{-n/2} \times \exp\{i \text{tr}[M(I - 2i B \Sigma / \sqrt{n})^{-1} B / \sqrt{n}]\} \quad (3.7)$$

where  $B = \sum_{i=1}^k t_i \text{diag}(c_{i1}, \dots, c_{ip})$ . Also,

$$\begin{aligned} E_2(t) &= E\left[\frac{i}{\sqrt{n}} \sum_{i=1}^k \sum_{\alpha=1}^r t_i a_{i\alpha} \text{tr} Z_{\alpha}^{(2)} \times \exp\left\{i \sum_{i_1=1}^k \sum_{\alpha_1=1}^r t_{i_1} a_{i_1 \alpha_1} \times \text{tr} Z_{\alpha_1}^{(1)}\right\}\right] \\ &= E_1(t) \frac{i}{\sqrt{n}} \sum_{i=1}^k \sum_{\alpha=1}^r \sum_{\beta \neq \alpha} \sum_{j_1 \in J_{\alpha}} \sum_{j_2 \in J_{\beta}} t_i a_{i\alpha} \theta_{\alpha\beta}^{-1} \\ &\quad \times \frac{1}{n} \left\{ \sum_{j=1}^n (\sigma_{j_1 j_1}^* \sigma_{j_2 j_2}^* + \sigma_{j_1 j_2}^{*2} + \sigma_{j_1 j_1}^* \xi_{j j_2}^2 + 2\sigma_{j_1 j_2}^* \xi_{j j_1} \xi_{j j_2} \right. \\ &\quad \left. + \sigma_{j_2 j_2}^* \xi_{j j_1}^2) \right. \\ &\quad \left. + \left[ \sum_{j=1}^n (\sigma_{j_1 j_2}^* + \xi_{j j_1} \xi_{j j_2}) \right]^2 \right\} \quad (3.8) \end{aligned}$$

$$\begin{aligned} E_3(t) &= E\left[\frac{i}{2\sqrt{n}} \sum_{i=1}^k \sum_{\alpha=1}^r \sum_{\beta=1}^r t_i a_{i\alpha\beta} (\text{tr} Z_{\alpha}^{(1)}) (\text{tr} Z_{\beta}^{(1)}) \right. \\ &\quad \left. \times \exp\left\{i \sum_{i_1=1}^k \sum_{\alpha_1=1}^r t_{i_1} a_{i_1 \alpha_1} \text{tr} Z_{\alpha_1}^{(1)}\right\}\right] \\ &= E_1(t) \frac{i}{2\sqrt{n}} \sum_{i=1}^k \sum_{\alpha=1}^r \sum_{\beta=1}^r \sum_{j_1 \in J_{\alpha}} \sum_{j_2 \in J_{\beta}} t_i a_{i\alpha\beta} \\ &\quad \times \left\{ \frac{1}{n} \sum_{j=1}^n (2\sigma_{j_1 j_2}^{*2} + 4\sigma_{j_1 j_2}^* \xi_{j j_1} \xi_{j j_2}) \right\} \end{aligned}$$

$$\begin{aligned}
& + \frac{1}{n} \sum_{j=1}^n \sum_{m=1}^n (\sigma_{j_1 j_1}^* + \xi_{j j_1}^*) (\sigma_{j_2 j_2}^* + \xi_{m j_2}^2) - \lambda_{j_1} \sum_{j=1}^n (\sigma_{j_2 j_2}^* + \xi_{j j_2}^2) \\
& - \lambda_{j_2} \sum_{j=1}^n (\sigma_{j_1 j_1}^* + \xi_{j j_1}^2) + n \lambda_{j_1} \lambda_{j_2} \} \quad (3.9)
\end{aligned}$$

where

$$\begin{aligned}
\Sigma^* &= \Sigma \left( I - \frac{2iB_1 \Sigma}{\sqrt{n}} \right)^{-1} = (\sigma_{ij}^*) \\
\xi_j &= \left( I - \frac{2i \Sigma B_1}{\sqrt{n}} \right)^{-1} \mu_j = (\xi_{j1}, \dots, \xi_{jp})' \quad (3.10)
\end{aligned}$$

Using the expansion that

$$|I-A|^{-\beta} = \exp \beta \left( \sum_{j=1}^{\infty} \frac{\text{tr } A^j}{j} \right)$$

in (3.7) and

$$(I-A)^{-1} = \sum_{j=0}^{\infty} A^j$$

in (3.10), we obtain

$$\sigma_{j_1 j_2}^* = \sigma_{j_1 j_2} + \frac{2i}{\sqrt{n}} \sum_{i=1}^k \sum_{j=1}^p t_i c_{ij} \sigma_{j_1 j} \sigma_{j_2 j} + O(n^{-1})$$

$$\xi_{j j_1} = \mu_{j j_1} + \frac{2i}{\sqrt{n}} \sum_{i=1}^k \sum_{m=1}^p t_i c_{im} \sigma_{j_1 m} \mu_{j m} + O(n^{-1})$$

Eq. (3.7), (3.8), (3.9) lead to

$$\begin{aligned}
\psi(\underline{t}) &= \exp\left(-\frac{1}{2} \underline{t}' Q \underline{t}\right) \\
&\times \left\{ 1 + \frac{1}{\sqrt{n}} \sum_{i=1}^k i t_i (d_1 + d_2) + \frac{1}{\sqrt{n}} \sum_{i_1 i_2 i_3}^k (i^3 t_{i_1} t_{i_2} t_{i_3}) (g_1 + g_2 + g_3) \right\} \\
&+ O(n^{-1}) \quad (3.11)
\end{aligned}$$

where  $Q = (Q_{i_1 i_2})$ ,  $Q_{i_1 i_2} = 2 \text{tr } R^{(i_1)} R^{(i_2)} + 4 \text{tr } R^{(i_1)} \psi^{(i_2)}$ ,

and  $Q$  is assumed to be nonsingular. Also,

$$d_1 = \sum_{\alpha=1}^r \sum_{\beta \neq \alpha}^r \sum_{j_1 \in J_\alpha} \sum_{j_2 \in J_\beta} a_{i\alpha} \theta_{\alpha\beta}^{-1} (\sigma_{j_1 j_1} \sigma_{j_2 j_2} + \sigma_{j_1 j_2}^2 + 2\sigma_{j_1 j_2} v_{j_1 j_2} + \sigma_{j_1 j_1} v_{j_2 j_2} + \sigma_{j_2 j_2} v_{j_1 j_1})$$

$$d_2 = \sum_{\alpha=1}^r \sum_{\beta=1}^r \sum_{j_1 \in J_\alpha} \sum_{j_2 \in J_\beta} a_{i\alpha\beta} (\sigma_{j_1 j_2}^2 + 2\sigma_{j_1 j_2} v_{j_1 j_2})$$

$$g_1 = \frac{4}{3} \text{tr } R^{(i_1)} R^{(i_2)} R^{(i_3)} + 4 \text{tr } R^{(i_1)} R^{(i_2)} \psi^{(i_3)}$$

$$g_2 = 4 \sum_{\alpha=1}^r \sum_{\beta \neq \alpha}^r \sum_{j_1 \in J_\alpha} \sum_{j_2 \in J_\beta} a_{i_1 \alpha} \theta_{\alpha\beta}^{-1} (\varepsilon_{j_1 j_2}^{(i_2)} + \tau_{j_1 j_2}^{(i_2)} + \tau_{j_2 j_1}^{(i_2)}) \times (\varepsilon_{j_1 j_2}^{(i_3)} + \tau_{j_1 j_2}^{(i_3)} + \tau_{j_2 j_1}^{(i_3)})$$

$$g_3 = 2 \sum_{\alpha=1}^r \sum_{\beta=1}^r \sum_{j_1 \in J_\alpha} \sum_{j_2 \in J_\beta} a_{i_1 \alpha\beta} (\varepsilon_{j_1 j_1}^{(i_2)} + 2\tau_{j_1 j_1}^{(i_2)}) (\varepsilon_{j_2 j_2}^{(i_3)} + 2\tau_{j_2 j_2}^{(i_3)})$$

(3.12)

We define here  $M/n = (v_{ij})$

$$C^{(i)} = \text{diag}(c_{i1}, \dots, c_{ip})$$

$$R^{(i)} = C^{(i)} \Sigma, \psi^{(i)} = C^{(i)} \frac{M}{n}, \varepsilon^{(i)} = \Sigma C^{(i)} \Sigma, \tau^{(i)} = \frac{M}{n} C^{(i)} \Sigma$$

(3.13)

where  $A_{ij}$  denotes the  $(i,j)$ th element of matrix  $A = (A_{ij})$ .

Now inverting (3.11) we obtain the following expansion for density of  $\underline{L} = (L_1, \dots, L_k)$

$$f(L_1, \dots, L_k) = N(\underline{L}, Q) \times \left[ 1 + \frac{1}{\sqrt{n}} \sum_{i=1}^k H_i(\underline{L})(d_1 + d_2) + \frac{1}{\sqrt{n}} \sum_{i_1, i_2, i_3}^k H_{i_1 i_2 i_3}(\underline{L})(g_1 + g_2 + g_3) \right] + O(n^{-1}) \quad (3.14)$$

where  $N(\underline{L}, Q) = \frac{1}{(2\pi)^{k/2} |Q|^{1/2}} \exp(-\frac{1}{2} \underline{L}' Q^{-1} \underline{L})$

$$H_{j_1, \dots, j_s}(\underline{L}) N(\underline{L}, Q) = (-1)^s \frac{\partial^s}{\partial L_{j_1} \dots \partial L_{j_s}} N(\underline{L}, Q) \quad (3.15)$$

Now, let  $T_i(\ell_1, \dots, \ell_p) = \ell_i$ . Then  $L_i = \sqrt{n} (\ell_i - \lambda_i)$ . Using Eq. (3.14), we obtain the following expression for the joint density of the roots  $\ell_1, \dots, \ell_p$  when  $\Sigma = \sigma^2 I$ :

$$f(L_1, \dots, L_p) = N(\underline{L}, Q) \left\{ 1 + \frac{1}{\sqrt{n}} \sum_{i=1}^p H_i(\underline{L}) \sum_{j \neq i} \theta_{ij}^{-1} (\sigma^2 \lambda_i + \sigma^2 \lambda_j - \sigma^4) + \frac{4\sigma^4}{\sqrt{n}} \sum_{i=1}^p H_{iii}(\underline{L}) (\lambda_i - \frac{2}{3} \sigma^2) \right\} + O(n^{-1}) \quad (3.16)$$

where  $Q = \text{diag}(Q_1, \dots, Q_p)$  and  $Q_i = 2\sigma^2(2\lambda_i - \sigma^2)$ . When  $\Sigma = \sigma^2 I$  and  $\lambda_1 > \dots > \lambda_t = \lambda_{t+1} = \dots = \lambda_p = \sigma^2$ , Carter and Srivastava (1979) obtained an alternative expression for the joint density of  $\ell_1, \dots, \ell_p$  by using a different method.

The general  $k$ -dimensional Hermite polynomial of order  $s \geq 0$  is denoted by  $H_{i_1, i_2, \dots, i_s}(\underline{x}; \Delta)$  where  $\underline{x} = (x_1, \dots, x_k)'$  is the polynomial variate and  $\Delta = (\delta_{ij})$  is a positive-definite  $k \times k$  matrix,  $0 \leq i_j \leq k$  for  $j=1, \dots, s$  and

$$H_{i_1, i_2, \dots, i_s}(\underline{x}, \Delta) N(\underline{x}, Q) = (-1)^s \frac{\partial^s}{\partial x_{i_1} \dots \partial x_{i_s}} N(\underline{x}, Q) \quad (3.17)$$

where

$$N(\underline{x}, Q) = \frac{1}{(2\pi)^{k/2} |Q|^{1/2}} \exp\left(-\frac{1}{2} \underline{x}' Q^{-1} \underline{x}\right) \quad (3.18)$$

and

$$\Delta = Q^{-1}$$

For dimension  $k=1$ ,  $Q=\tau^2$  is a scalar and  $\Delta=\delta=\frac{1}{\tau^2}$

$$H_1(x, \delta) = x\delta$$

$$H_{111}(x, \Delta) = x^3 \delta^3 - 3x\delta^2 \quad (3.19)$$

and

$$\int_{-\infty}^{\tau a} H_1(x, \Delta) N(x, Q) dx = -\frac{1}{\tau} \frac{1}{\sqrt{2\pi}} \exp\left(\frac{-a^2}{2}\right) \quad (3.20)$$

$$\int_{-\infty}^{\tau a} H_{111}(x, \Delta) N(x, Q) dx = \frac{1}{\tau^3} (1-a^2) \frac{1}{\sqrt{2\pi}} \exp\left(\frac{-a^2}{2}\right)$$

For the dimension  $k=2$ , let  $Q = \begin{pmatrix} \tau_1^2 & \rho\tau_1\tau_2 \\ \rho\tau_1\tau_2 & \tau_2^2 \end{pmatrix}$ . Then

$$\Delta = Q^{-1} = \frac{1}{\tau_1^2 \tau_2^2 - (\rho\tau_1\tau_2)^2} \begin{pmatrix} \tau_2^2 & -\rho\tau_1\tau_2 \\ -\rho\tau_1\tau_2 & \tau_1^2 \end{pmatrix} = \begin{pmatrix} \delta_{11} & \delta_{12} \\ \delta_{21} & \delta_{22} \end{pmatrix} \quad (3.21)$$

Also, we have

$$\begin{aligned}
 H_1(\underline{x}, \Delta) &= x_1 \delta_{11} + x_2 \delta_{12} \\
 H_{111}(\underline{x}, \Delta) &= x_1^3 \delta_{11}^3 + 3 x_1^2 x_2 \delta_{11}^2 \delta_{12} + 3 x_1 x_2^2 \delta_{11} \delta_{12}^2 \\
 &\quad + x_2^3 \delta_{12}^3 - 3 x_1 \delta_{11}^2 - 3 x_2 \delta_{11} \delta_{12} \quad (3.22)
 \end{aligned}$$

$$\begin{aligned}
 H_{112}(\underline{x}, \Delta) &= x_1^3 \delta_{11}^2 \delta_{12} + x_1^2 x_2 (2 \delta_{11} \delta_{12}^2 + \delta_{11}^2 \delta_{22}) \\
 &\quad + x_1 x_2^2 (\delta_{12}^3 + 2 \delta_{11} \delta_{12} \delta_{22}) + x_2^3 \delta_{12}^2 \delta_{22} \\
 &\quad - 3 x_1 \delta_{11} \delta_{12} - x_2 (2 \delta_{12}^2 + \delta_{11} \delta_{22}).
 \end{aligned}$$

Similar equations for  $H_2(\underline{x}, \Delta)$ ,  $H_{222}(\underline{x}, \Delta)$  and  $H_{122}(\underline{x}, \Delta)$  are obtained by interchanging subscript 1, 2 in  $H_1(\underline{x}, \Delta)$ ,  $H_{111}(\underline{x}, \Delta)$  and  $H_{112}(\underline{x}, \Delta)$  respectively.

Now

$$\int_{-\infty}^{\tau_2^b} \int_{-\infty}^{\tau_1^a} x_1^r x_2^s N(\underline{x}, Q) dx_1 dx_2 = \int_{-\infty}^b \int_{-\infty}^a \tau_1^r x_1^r \tau_2^s x_2^s N(\underline{x}, R) dx_1 dx_2 \quad (3.23)$$

by changing of variables and  $R = \begin{pmatrix} 1 & \rho \\ \rho & 1 \end{pmatrix}$  is the correlation matrix.

Define

$$\mu_{r,s} = \int_{-\infty}^b \int_{-\infty}^a x_1^r x_2^s N(\underline{x}, R) dx_1 dx_2 \quad (3.24)$$

Integrating by parts, we have

$$\mu_{1,0} = -(\phi(a)\phi(B) + \rho\phi(b)\phi(A)) \quad (3.25)$$

$$\mu_{1,1} = -\rho b\phi(b)\phi(A) - \rho a\phi(a)\phi(B) + (1-\rho^2) N(a,b;R)$$

$$+ \rho \mu_{0,0}$$

$$\begin{aligned} \mu_{2,0} &= \mu_{0,0} - a \phi(a) \phi(B) - \rho^2 b \phi(b) \phi(A) \\ &+ \rho(1-\rho^2) N(a,b;R) \end{aligned}$$

In general, the recursive relation is

$$\begin{aligned} \mu_{r,s+1} - b \mu_{r,s} &= (1-\rho^2)_s \mu_{r,s-1} + \mu_{r+1,s} \rho \\ &- (1-\rho^2) b(s-1) \mu_{r,s-2} - b \rho \mu_{r+1,s} \end{aligned} \quad (3.26)$$

$$\begin{aligned} \mu_{r+1,s} - a \mu_{r,s} &= (1-\rho^2)_r \mu_{r-1,s} + \mu_{r,s+1} \rho - (1-\rho^2) a (r-1) \\ &\times \mu_{r-2,s} - a \rho \mu_{r-1,s+1} \end{aligned}$$

and

$$\begin{aligned} \mu_{3,0} &= -(a^2+2) \phi(a) \phi(B) + \rho(\rho^2 - \rho^2 b^2 - 3) \phi(b) \phi(A) \\ &+ (a+\rho b) \rho(1-\rho^2) N(a,b;R) \end{aligned} \quad (3.27)$$

$$\begin{aligned} \mu_{2,1} &= -\rho(a^2+2) \phi(a) \phi(B) - (1 + \rho^2 + \rho^2 b^2) \phi(b) \phi(A) \\ &+ (1-\rho^2)(a+\rho b) N(a,b;R) \end{aligned}$$

where

$$A = (1-\rho^2)^{-\frac{1}{2}} (a-\rho b), \quad B = (1-\rho^2)^{-\frac{1}{2}} (b-\rho a)$$

$$\phi(a) = \frac{1}{\sqrt{2\pi}} \exp \frac{-a^2}{2} \quad (3.28)$$

$$\Phi(A) = \int_{-\infty}^A \phi(t) dt$$

$\mu_{0,1}$ ,  $\mu_{0,2}$ ,  $\mu_{0,3}$  and  $\mu_{1,2}$  are obtained by interchanging  $a$  with  $b$  and  $A$  with  $B$  in  $\mu_{1,0}$ ,  $\mu_{2,0}$ ,  $\mu_{3,0}$  and  $\mu_{2,1}$  respectively.

#### 4. Applications in Investigation of the Structures of Interactions

In this section, we discuss some applications of the results of Section 3 in studying the power functions of various tests for the hypothesis of no interaction in two-way classification model with one observation per cell.

Consider the model

$$x_{ij} = \mu + \alpha_i + \beta_j + \eta_{ij} + \epsilon_{ij} \quad (4.1)$$

for  $i = 1, \dots, u$ ,  $j = 1, 2, \dots, s$ . Here  $x_{ij}$  denotes the observation in  $i$ -th row and  $j$ -th column,  $\mu$  is the general mean,  $\alpha_i$  denotes the effect due to  $i$ -th row,  $\beta_j$  denotes the effect due to  $j$ -th column and  $\eta_{ij}$  denotes the interaction of  $i$ -th row and  $j$ -th column. Also, we assume that  $\epsilon_{ij}$ 's are distributed independently and normally with mean 0 and variance  $\sigma^2$ . Now, let  $d_{ij} = x_{ij} - \bar{x}_{i.} - \bar{x}_{.j} + \bar{x}_{..}$ , where  $s\bar{x}_{i.} = \sum_{j=1}^s x_{ij}$ ,  $u\bar{x}_{.j} = \sum_{i=1}^u x_{ij}$  and  $us\bar{x}_{..} = \sum_{i=1}^u \sum_{j=1}^s x_{ij}$ . Also, let  $D = (d_{ij})$ ,  $X = (x_{ij})$ ,  $W = C_u' X C_s C_s' X' C_u$  where  $C_u$  is chosen such that  $C_u' C_u = I_{u-1}$  and  $C_u C_u' = I_u - \frac{1}{u} J_u$  where  $J_u$  is the  $u \times u$  matrix with all its elements equal to unity. The non-zero eigenvalues of  $DD'$  are the same (e.g., see Johnson and Graybill (1972)) as the nonzero eigenvalues of  $W$ . Also, the columns of  $C_u' X$  are distributed independently as multivariate normal with mean vectors given by  $E(C_u' X) = C_u' M_0$  and a common covariance matrix  $\sigma^2 I_{u-1}$  where  $M_0 = (m_{ij})$  and  $m_{ij} = \mu + \alpha_i + \beta_j + \eta_{ij}$ . In addition,

$$E(W/(s-1)) = \sigma^2 I_{u-1} + \{C_u' M_o C_s C_s' M_o C_u / (s-1)\} \\ = \Sigma_0 \quad (4.2)$$

and  $C_u' M_o C_s C_s' M_o C_u = C_u' \eta \eta' C_u = \Omega$  where  $\eta = (\eta_{ij})$ . So,  $W$  is distributed as the noncentral Wishart matrix with  $(s-1)$  degrees of freedom and noncentrality matrix  $\Omega$ . When  $\eta=0$ ,  $W$  is distributed as the central Wishart matrix with  $(s-1)$  degrees of freedom.

Let  $\lambda_1 \geq \dots \geq \lambda_{u-1}$  be the eigenvalues of  $W/(s-1)$  and let  $\lambda_1 \geq \dots \geq \lambda_{u-1}$  be the nonzero roots of  $\Sigma_0$ . Then, the problem of testing the hypothesis  $H: \Omega = 0$  is equivalent to testing the hypothesis that the eigenvalues of  $\Sigma_0$  are equal. Suppose  $\eta = \lambda \alpha \beta'$  where  $\alpha' = (\alpha_1, \dots, \alpha_u)$  and  $\beta' = (\beta_1, \dots, \beta_s)$ . Then  $\Omega = \lambda^2 C_u' \alpha \beta' \beta \alpha' C_u$ , and the nonzero root of  $\Omega$  is  $\lambda \beta' \beta \alpha' \alpha$ . Next, we will assume that the rank of  $\eta$  is  $c$ . Then, using the well-known singular value decomposition of the matrix, we can write  $\eta$  as

$$\eta = \lambda_1 w_1 v_1' + \dots + \lambda_c w_c v_c' \quad (4.3)$$

The nonzero eigenvalues of  $\eta \eta'$  are  $\lambda_1^2, \dots, \lambda_c^2$  and the associated eigenvectors are  $w_1, \dots, w_c$ . The eigenvectors of  $\eta' \eta$  corresponding to the eigenvalues  $\lambda_1^2, \dots, \lambda_c^2$  are  $v_1, \dots, v_c$ . The nonzero eigenvalues of  $\Omega$  are  $\lambda_1^2, \dots, \lambda_c^2$ .

The problem of testing the hypothesis of no interaction in two-way classification with one observation per cell was studied by several authors (e.g., see Tukey (1949)

and Williams (1951)). The statistic proposed by Tukey is given by  $(\hat{\alpha}'\hat{\eta}\hat{\beta})^2/(\hat{\alpha}'\hat{\alpha})(\hat{\beta}'\hat{\beta})$  where  $\hat{\alpha}' = (\hat{\alpha}_1, \dots, \hat{\alpha}_u)$ ,  $\hat{\beta}' = (\hat{\beta}_1, \dots, \hat{\beta}_s)$ ,  $\hat{\eta} = (\hat{\eta}_{ij})$ ,  $\hat{\alpha}_i = \bar{x}_{i.} - \bar{x}_{..}$ ,  $\hat{\beta}_j = \bar{x}_{.j} - \bar{x}_{..}$  and  $\hat{\eta}_{ij} = x_{ij} - \bar{x}_{i.} - \bar{x}_{.j} + \bar{x}_{..}$ . Gollob (1968) and Mandel (1969) considered the problem of testing the hypotheses  $\lambda_j = 0$  individually under the model (4.3) by using the statistics  $F_j = \ell_j / (\ell_1 + \dots + \ell_{u-1})$ . Gollob treated  $\ell_1, \dots, \ell_{u-1}$  as independent chi-square variables to get an approximation to the distribution of  $F_j$ . But  $\ell_i$ 's are neither independent nor distributed as chi-square variables. Corsten and Van Eijnsbergen (1972) showed that the likelihood ratio statistic for testing the hypothesis  $\lambda_1 = \dots = \lambda_c = 0$  under the model (4.3) is  $(\ell_1 + \dots + \ell_c) / (\ell_{c+1} + \dots + \ell_{u-1})$ . When  $c=1$ , this statistic was derived independently by Johnson and Graybill (1972). Schuurmann, Krishnaiah and Chattopadhyay (1973) and Krishnaiah and Schuurmann (1974) discussed the problem of testing the hypotheses  $\lambda_i = 0$  simultaneously by applying the simultaneous tests of Krishnaiah and Waikar (1971) for the equality of the eigenvalues of the covariance matrix of the multivariate normal population. Ghosh and Sharma (1963) studied the power function of Tukey's test for  $\eta_{ij} = 0$  against the alternative that  $\eta_{ij} = \lambda \alpha_i \beta_j$ . Yochmowitz and Cornell (1978) derived the likelihood ratio test for  $\lambda_1 = \dots = \lambda_a = 0$  against the alternative that  $\lambda_a \neq 0$  and  $\lambda_{a+1} = \dots = \lambda_c = 0$ . We now compare the power functions of various procedures for

testing the hypothesis of no interaction.

Let  $T_1 = \ell_1 / \ell_{u-1}$ ,  $T_2 = (\text{tr } W / u - 1)^{u-1} / |W|$ ,  $T_3 = (u - c - 1) \times \ell_1 / (\ell_{c+1} + \dots + \ell_{u-1})$ ,  $T_4 = (u - c - 1)(\ell_1 + \dots + \ell_c) / c(\ell_{c+1} + \dots + \ell_{u-1})$ . When  $\sigma^2$  is unknown, we can use any of the above statistics for testing the hypothesis of no interaction.

If we use  $T_i$ , we accept or reject  $H_0$  according as

$$T_i \lesseqgtr c_\alpha \quad (4.4)$$

where

$$P[T_i \leq c_\alpha | H_0] = (1 - \alpha) \quad (4.5)$$

The test statistics  $T_1$  is based upon the statistic considered by Krishnaiah and Waikar (1971) for testing the sphericity, whereas the test statistic  $T_2$  is based upon the likelihood ratio test statistic for sphericity. The statistic  $T_4$  is the likelihood ratio test statistic (see Corsten and Van Eijnsbergen (1972)) for testing the hypothesis of no interaction of multiplicative components model (4.3).

Table 1 gives a comparison of the power functions of various procedures for testing the hypothesis of no interaction when  $\sigma^2$  is known. The rows corresponding to S denote the simulated values. The multivariate normal deviates are generated by the IMSL subroutine GGNRM, and 10,000 trials are performed for each case, the 95% confidence limit for each value is then  $1.96\{\hat{p}(1-\hat{p})/10,000\}^{1/2}$ , where  $\hat{p}$  is the actual value from the empirical trials.

The rows corresponding to  $N$  denote the values corresponding to the first term in the asymptotic expansion. The rows corresponding to  $N + O(n^{-\frac{1}{2}})$  give the values corresponding to the first two terms of the expansion.

TABLE 1  
Comparison of the Power Functions of the Tests  
for no Interaction When  $\sigma^2$  is Unknown

$p = 3, \alpha = 0.05$

$n$	$(\lambda_1, \lambda_2, \lambda_3)$	Type of Approximation	$\lambda_1/\lambda_p$	$\frac{(\text{tr}W/p)p}{ W }$	$(p-1)\lambda_1/(\lambda_2+\lambda_3+\dots+\lambda_p)$	$(p-2)(\lambda_1+\lambda_2)/2\lambda_3$
10	(12, 6, 1)	N	0.57	0.49		0.59
		$N+O(n^{-\frac{1}{2}})$	0.83	0.78		0.83
		S	0.81	0.77		0.82
10	(12, 3, 1)	N	0.57	0.63		0.44
		$N+O(n^{-\frac{1}{2}})$	0.85	0.90		0.71
		S	0.82	0.87		0.73
10	(12, 10, 1)	N	0.57	0.55		0.72
		$N+O(n^{-\frac{1}{2}})$	0.88	0.83		0.95
10	(7, 1, 1)	N			0.79	
		$N+O(n^{-\frac{1}{2}})$			0.95	
10	(12, 12, 1)	N				0.76
		$N+O(n^{-\frac{1}{2}})$				0.99
25	(4, 3.5, 1)	N	0.51	0.54		0.68
		$N+O(n^{-\frac{1}{2}})$	0.82	0.79		0.85
		S	0.79	0.77		0.84
25	(4, 1, 1)	N			0.89	
		$N+O(n^{-\frac{1}{2}})$			0.98	
25	(4, 4, 1)	N				0.74
		$N+O(n^{-\frac{1}{2}})$				0.90
		S				0.89

TABLE 1 (continued)

n	$(\lambda_1, \lambda_2, \lambda_3)$	Type of Approximation	$\ell_1 / \ell_p$	$\frac{(\text{tr}W/p)^p}{ W }$	$(p-1)\ell_1 / (\ell_2 + \ell_3)$	$(p-2)(\ell_1 + \ell_2) / 2\ell_3$
50	(2.5, 1.7, 1)	N N+O( $n^{-\frac{1}{2}}$ ) S	0.46 0.66 0.68	0.42 0.68 0.68		0.45 0.62 0.64
50	(2.5, 1, 1)	N N+O( $n^{-\frac{1}{2}}$ ) S			0.83 0.92 0.92	
50	(2.5, 2.5, 1)	N N+O( $n^{-\frac{1}{2}}$ )				0.72 0.86
100	(2, 1.5, 1)	N N+O( $n^{-\frac{1}{2}}$ ) S	0.57 0.75 0.75	0.54 0.77 0.76		0.57 0.72 0.73
100	(2, 1.15, 1)	N N+O( $n^{-\frac{1}{2}}$ ) S	0.57 0.82 0.82	0.65 0.86 0.85		0.32 0.60 0.63
100	(2, 1, 1)	N N+O( $n^{-\frac{1}{2}}$ )			0.88 0.95	
100	(2, 2, 1)	N N+O( $n^{-\frac{1}{2}}$ )				0.82 0.92

TABLE 1 (continued)  
 $p = 4, \alpha = 0.05$

$n$	$(\lambda_1, \lambda_2, \lambda_3, \lambda_4)$	Type of Approximation	$\ell_1/\ell_p$	$\frac{(\text{tr}W/p)^p}{ W }$	$\frac{(p-1)\ell_1}{\ell_2+\ell_3+\ell_4}$	$\frac{(p-1)\ell_1}{\ell_3+\ell_4}$	$\frac{(p-2)(\ell_1+\ell_2)}{2(\ell_3+\ell_4)}$	$\frac{(p-3)(\ell_1+\ell_2+\ell_3)}{3\ell_4}$
100	(3, 2.5, 2, 1)	N $N+O(n^{-\frac{1}{2}})$	0.45 0.78	0.36 0.76				0.59 0.80
		S	0.79	0.76				0.81
100	(3, 2.5, 1, 1)	N $N+O(n^{-\frac{1}{2}})$			0.80 1.00	0.90 1.00		
100	(3, 2.5, 2.5, 1)	N $N+O(n^{-\frac{1}{2}})$						0.70 0.88
100	(3, 1, 1, 1)	N $N+O(n^{-\frac{1}{2}})$			0.95 1.00			
100	(3, 3, 2, 1)	N $N+O(n^{-\frac{1}{2}})$						0.70 0.88
100	(3, 3, 3, 1)	N $N+O(n^{-\frac{1}{2}})$ S						0.83 0.98 0.96
100	(3, 3, 1, 1)	N $N+O(n^{-\frac{1}{2}})$					0.96 1.00	

TABLE 1 (continued)

 $p = 4, \alpha = 0.05$ 

$n$	$(\lambda_1, \lambda_2, \lambda_3, \lambda_4)$	Type of Approximation	$\ell_1 / \ell_p$	$\frac{(\text{tr} W / p)^p}{ W }$	$\frac{(p-1)\ell_1}{\ell_2 + \ell_3 + \ell_4}$	$\frac{(p-1)\ell_1}{\ell_3 + \ell_4}$	$\frac{(p-2)(\ell_1 + \ell_2)}{2(\ell_3 + \ell_4)}$	$\frac{(p-3)(\ell_1 + \ell_2 + \ell_3)}{3\ell_4}$
100	(3, 2, 5, 2, 1)	N	0.45	0.36				0.59
		$N+O(n^{-\frac{1}{2}})$	0.78	0.76				0.80
		S	0.79	0.76				0.81
100	(3, 2, 5, 1, 1)	N				0.80	0.90	
		$N+O(n^{-\frac{1}{2}})$				1.00	1.00	
100	(3, 2, 5, 2, 5, 1)	N						0.70
		$N+O(n^{-\frac{1}{2}})$						0.88
100	(3, 1, 1, 1)	N			0.95			
		$N+O(n^{-\frac{1}{2}})$			1.00			
100	(3, 3, 2, 1)	N						0.70
		$N+O(n^{-\frac{1}{2}})$						0.88
100	(3, 3, 3, 1)	N						0.83
		$N+O(n^{-\frac{1}{2}})$						0.98
		S						0.96
100	(3, 3, 1, 1)	N					0.96	
		$N+O(n^{-\frac{1}{2}})$					1.00	

The radius of convergence is

$$\gamma_0 = \frac{d}{2||V||} \quad (4.6)$$

where

$$V = \sqrt{n} (S/n - M)$$

We will choose  $n$  such that  $1/\sqrt{n} < \gamma_0$ . Now  $||V|| = CL(\sqrt{n}(S/n - M))$ , where  $CL(\cdot)$  denotes the largest root of  $\sqrt{n}((S/n) - M)$ , and is approximately distributed with mean 0 and variance  $2\sigma^2(\lambda_1 - \sigma^2)$ .

In Table 1, consider the entry when  $p=3$ ,  $\lambda_1=12$ ,  $\lambda_2 = 6$  and  $\lambda_3 = 1$ . In this case,  $\sigma^2 = 1$ ,  $d = \lambda_2 - \lambda_3 = 5$ , with one standard deviation,

$$\frac{d}{2\sqrt{2\sigma^2(2\lambda_1 - \sigma^2)}} \sim \frac{1}{\sqrt{7.5}}$$

where  $n=10$  is chosen.

When the entry is  $p = 4$ ,  $\lambda_1 = 2.5$ ,  $\lambda_2 = 1.7$  and  $\lambda_3 = 1$ , then  $\sigma^2 = 1$ ,  $\lambda_1 = 2.5$ ,  $d = \lambda_1 - \lambda_2 = 0.8$  and  $n = 50$ .

The table reveals that results based on normal approximations are not sufficiently accurate for  $n$  as large as 100, while the asymptotic expression taking the term of order  $n^{-\frac{1}{2}}$  achieves numerical accuracy for moderate sample sizes. This suggests that care should be given for the statistical inferences which are based on the normal approximations.

Next, consider the model (4.1) when  $\sigma^2$  is known. In this case, we accept or reject the hypothesis  $\lambda_1 = \dots = \lambda_c = 0$  according as

$$\frac{\ell_1}{\sigma^2} \lessgtr d_{2\alpha} \quad (4.7)$$

where

$$P \left[ \frac{\ell_1}{\sigma^2} \leq d_{2\alpha} | H \right] = (1-\alpha) \quad (4.8)$$

When  $H$  is rejected, the hypothesis  $\theta_i = 0$  is accepted or rejected according as  $(\ell_i/\sigma^2) \leq d_{2\alpha}$ . When  $H$  is true,  $\ell_1$  is the largest eigenvalue of the central Wishart matrix. Exact distribution of this statistic is given in Krishnaiah and Chang (1971) and exact percentage points are given in Krishnaiah (1980). When  $H$  is not true, an asymptotic expression for the distribution of  $\ell_1$  can be obtained as a special case of (3.14) if  $\lambda_1$  is different from  $\theta_2, \dots, \theta_r$ .

When  $\lambda_1 > \dots > \lambda_2 > 0$ , Srivastava and Carter (1980) obtained asymptotic expression of  $\log(\ell_1/\ell_1 + \dots + \ell_{u-1})$  and  $\tau_2^{1/(u-1)}$  by a different method. For a review of the literature on tests for no interaction in two way classification model with one observation per cell, the reader is referred to Krishnaiah and Yochmowitz (1980).

## 5. Applications in Cluster Analysis and Reduction of Dimensionality

Let  $X_1, \dots, X_N$  be independent  $p$ -dimensional random variables. We consider using the sample covariance matrix

$$S = \sum_{i=1}^N (\underline{X}_i - \bar{\underline{X}})(\underline{X}_i - \bar{\underline{X}})'$$

where  $\bar{\underline{X}} = N^{-1}(\underline{X}_1 + \dots + \underline{X}_N)$ . We wish to test the hypothesis that  $\underline{X}_i$ 's come from a single multivariate normal population with covariance  $\Sigma$  against the hypothesis that they come from a mixture of  $k \leq p$  such populations, differing in means. We assume  $\Sigma = \sigma^2 I$ . For  $k=2$  the null hypothesis  $H_1$  and the alternative hypothesis  $H_2$  are given by

$$H_1: \underline{X}_i \sim N(\underline{\mu}, \sigma^2 I)$$

$$H_2: \underline{X}_i \sim \pi N(\underline{\mu}_1, \sigma^2 I) + (1-\pi) N(\underline{\mu}_2, \sigma^2 I)$$

where  $\pi$  is the mixing probability. Under  $H_2$  it is known (e.g. see Bryant (1975)) that

$$S \sim \sum_{j=0}^N \binom{N}{j} \pi^j (1-\pi)^{N-j} W_p(N-1, \sigma^2 I, M_j)$$

$$M_j = N^{-1} j(N-j)(\underline{\mu}_1 - \underline{\mu}_2)(\underline{\mu}_1 - \underline{\mu}_2)'$$

$M_j$  is of rank 1. Now, let  $\Delta^2 = (\underline{\mu}_1 - \underline{\mu}_2)'(\underline{\mu}_1 - \underline{\mu}_2)/\sigma^2$  which is proportional to the largest root of  $M_j/(N-1)$ . When the null hypothesis is true, we know that

$$S \sim W_p(N-1, \sigma^2 I, 0).$$

Let the test statistics  $T_1$  and  $T_2$  be given as below:

$$T_1 = \lambda_1 / \sigma^2 \quad (5.1)$$

$$T_2 = \frac{(p-1)\lambda_1}{\lambda_2 + \dots + \lambda_p} \quad (5.2)$$

where  $\lambda_1 \geq \dots \geq \lambda_p$  are eigenvalues of  $S/(N-1)$ .

If we use the statistics  $T_i$  to test  $H_1$ , then we accept or reject  $H_1$  according as

$$T_i \leq c_{\alpha i} \quad (5.3)$$

where

$$P\{T_i \leq c_{\alpha i} | H_1\} = 1 - \alpha \quad (5.4)$$

Let  $f_j(\cdot)$  be the asymptotic density of a function of the eigenvalues of  $S/(N-1)$  when  $j$  of the samples come from population 1. Under this condition  $S \sim W_p(N-1, \sigma^2 I, M_j)$ . So the unconditional asymptotic density function of the function of eigenvalue of  $S$  is

$$\sum_{j=0}^N \binom{N}{j} \pi^j (1-\pi)^{N-j} f_j(\cdot)$$

The following table gives a comparison of the asymptotic power value with the simulated value of tests of  $H_1$  against  $H_2$  for  $p=4$ ,  $\alpha=0.05$ ,  $\pi$  = mixing probability,  $\Delta = ||\mu_1 - \mu_2||/\sigma$ ,  $N$  = sample size.

Test	$\pi = .25$			$\pi = .50$		
	$\Delta=1$	2	3	$\Delta=1$	2	3
$T_1$	.02	.42	.95	.02	.63	1.00
Simu.	.07	.45	.94	.07	.62	1.00

N= 51

		$\pi = .25$			$\pi = .50$		
Test		$\Delta=1$	2	3	$\Delta=1$	2	3
N=51	$T_2$	.07	.68	.99	.11	.88	1.00
	Simu.	.11	.70	.98	.14	.86	1.00
	$T_1$	.05	.76	1.00	.08	.96	1.00
	Simu.	.13	.78	1.00	.21	.96	1.00
N=101	$T_2$	.13	.93	1.00	.22	1.00	1.00
	Simu.	.13	.93	1.00	.27	1.00	1.00

When  $\Delta = 1$ , the largest roots of  $M_j/(N-1)$  are close to zero, the radius of convergence for perturbation approximation of eigenvalues is small, and the asymptotic expression does not give good approximation. Note, that if the means under  $H_2$  are separated by more than two or three standard derivations, that is, for  $\Delta = 2, 3$ , one may reasonably expect to detect the presence of two components, while if they are separated by less than two standard derivations the detection generally will not be good.

Consider  $k$   $p$ -variate normal populations with unknown mean vectors  $\mu_1, \dots, \mu_k$  and a common known covariance matrix  $\Sigma$ . We assume that  $n_i$  observations are available from  $i$ -th population and the sample mean vector and corrected sum of squares and cross-products (SP) matrix based on these observations are respectively given by  $\bar{X}_i$  and  $S_i$ . The SP matrix which explain the variation between groups is given by

$$B = \sum_{i=1}^k n_i (\bar{X}_i - \bar{X}_{..})(\bar{X}_i - \bar{X}_{..})'$$

$$n_i \bar{X}_{i.} = \sum_{j=1}^{n_i} X_{ij}, \quad n \bar{X}_{..} = \sum_{i=1}^k \sum_{j=1}^{n_i} X_{ij} \quad (5.5)$$

and  $n = n_1 + \dots + n_k$ . Now, let  $\lambda_1 \geq \dots \geq \lambda_p$  denote the eigenvalues of  $B\Sigma^{-1}$ . Fisher (1939) studied the problem of whether there are any structural relations between the  $p$  components of mean vectors. This is equivalent to testing the hypothesis  $H_0$  where

$$H_0 : C\mu_i = \xi \quad (5.6)$$

for  $i = 1, 2, \dots, k$  where  $C: s \times p$  and  $\xi$  are unknown and the rank of  $C$  is  $s$ . We assume that  $t < k-1$  where  $t = p-s$ . The likelihood ratio statistic for testing  $H_0$  is given by  $U_1 = (\lambda_{t+1} + \dots + \lambda_p)$ . A detailed discussion of the above procedure was given in Rao (1965). The statistic  $B\Sigma^{-1}$  is distributed as the noncentral Wishart matrix with  $\gamma$  degrees of freedom and

$$E(B\Sigma^{-1}) = (k-1)I + \Omega\Sigma^{-1} = \Sigma_* \quad (5.7)$$

where  $\gamma = k-1$

$$\Omega = \sum_{i=1}^k n_i (\mu_i - \bar{\mu})(\mu_i - \bar{\mu})' \quad (5.8)$$

and  $n\bar{\mu} = n_1\mu_1 + \dots + n_k\mu_k$ . Also, let  $\lambda_1 \geq \dots \geq \lambda_p$  denote the roots of  $\Sigma_*$ . The distribution of  $U_1$  can be obtained as a special case of (3.14) and so the power of the likelihood ratio test for  $H_0$  can be studied using the results in this paper.

## 6. Asymptotic Joint Distribution of Functions of the Eigenvalues of Multivariate Quadratic Form

In this section we shall derive the joint asymptotic distributions of functions of eigenvalues of the multivariate quadratic form  $S = XGX'$ , where we assume that  $G^S = O(1)$ , for whatever power  $s$  raised on  $G$ .  $G$  is a symmetric matrix and the columns of  $X: p \times n$  are distributed independently as multivariate normal with covariance matrix  $\Sigma = (\sigma_{ij})$  and means  $E(X) = U = (\mu_1, \dots, \mu_n)$ . Then

$$E(S/n) = \frac{\text{tr}G}{n} \Sigma + \frac{UGU'}{n} = \Lambda = \text{diag}(\lambda_1, \dots, \lambda_p), \quad (6.1)$$

Also assume that

$$\lambda_{q_1 + \dots + q_{\alpha-1} + 1} = \dots = \lambda_{q_1 + \dots + q_{\alpha}} = \theta_{\alpha} \quad (6.2)$$

for  $\alpha=1, 2, \dots, r$ ,  $q_1 + \dots + q_r = p$ , and  $q_0=0$  and let  $\ell_1 \geq \dots \geq \ell_p$  be the eigenvalues of  $S/n$ .

We consider the joint asymptotic distribution of  $L_1, \dots, L_k$  where  $L_j = \sqrt{n} \{T_j(\ell_1, \dots, \ell_p) - T_j(\lambda_1, \dots, \lambda_p)\}$ , which satisfy assumptions of (3.2). Let

$$S/n = \Lambda + V/\sqrt{n} \quad (6.3)$$

Using the Taylor expansion of  $T_i(\ell_1, \dots, \ell_p)$  for  $i=1, 2, \dots, k$ , as in (3.3) and the application of Lemma 2.1 we obtain the same equations as (3.5), (3.6) and the characteristic function

$$\Psi(t_1, \dots, t_k) = E_1(t) + E_2(t) + E_3(t) + O(n^{-1})$$

$$E_1(t) = \exp(-i \sqrt{n} \operatorname{tr} B_1 \Lambda) |I - 2i(G \otimes B_1 \Sigma / \sqrt{n})|^{-\frac{1}{2}}$$

$$\exp[i \operatorname{tr} \xi \xi' (I - 2i G \otimes B_1 \Sigma / \sqrt{n})^{-1} G \otimes B_1 / \sqrt{n}]$$

$$\begin{aligned} E_2(t) = E_1(t) & \frac{1}{\sqrt{n}} \sum_{i=1}^k \sum_{\alpha=1}^r \sum_{\beta \neq \alpha} \sum_{j_1 \in J_\alpha} \sum_{j_2 \in J_\beta} t_i a_{i\alpha} \theta_{\alpha\beta}^{-1} \\ & \times \frac{1}{n} \sum_{k_1, k_2, k_3, k_4}^n g_{k_1 k_2} g_{k_3 k_4} (\sigma_{i_1 i_2}^* \sigma_{i_3 i_4}^* \\ & + \sigma_{i_1 i_3}^* \sigma_{i_2 i_4}^* + \sigma_{i_1 i_4}^* \sigma_{i_2 i_3}^* \\ & + \sigma_{i_1 i_2}^* \pi_{i_3} \pi_{i_4} + \sigma_{i_1 i_3}^* \pi_{i_2} \pi_{i_4} + \sigma_{i_1 i_4}^* \pi_{i_2} \pi_{i_3} + \sigma_{i_2 i_3}^* \pi_{i_1} \pi_{i_4} \\ & + \sigma_{i_2 i_4}^* \pi_{i_1} \pi_{i_3} + \sigma_{i_3 i_4}^* \pi_{i_1} \pi_{i_2} + \pi_{i_1} \pi_{i_2} \pi_{i_3} \pi_{i_4}) \end{aligned} \quad (6.4)$$

$$\begin{aligned} E_3(t) = E_1(t) & \frac{1}{2\sqrt{n}} \sum_{i=1}^k \sum_{\alpha=1}^r \sum_{\beta=1}^r \sum_{j_1 \in J_\alpha} \sum_{j_2 \in J_\beta} t_i a_{i\alpha\beta} \\ & \times \left[ \frac{1}{n} \sum_{k_1, k_2, k_3, k_4}^n g_{k_1 k_2} g_{k_3 k_4} (\sigma_{m_1 m_2}^* \sigma_{m_3 m_4}^* + \sigma_{m_1 m_3}^* \sigma_{m_2 m_4}^* \right. \\ & + \sigma_{m_1 m_4}^* \sigma_{m_2 m_3}^* \\ & + \sigma_{m_1 m_2}^* \pi_{m_3} \pi_{m_4} + \sigma_{m_1 m_3}^* \pi_{m_2} \pi_{m_4} + \sigma_{m_1 m_4}^* \pi_{m_2} \pi_{m_3} \\ & + \sigma_{m_2 m_3}^* \pi_{m_1} \pi_{m_4} \\ & + \sigma_{m_2 m_4}^* \pi_{m_1} \pi_{m_3} + \sigma_{m_3 m_4}^* \pi_{m_1} \pi_{m_2} + \pi_{m_1} \pi_{m_2} \pi_{m_3} \pi_{m_4}) \\ & \left. - \lambda_{j_2} \sum_{k_1, k_2} g_{k_1 k_2} (\sigma_{m_1 m_2}^* + \pi_{m_1} \pi_{m_2}) - \lambda_{j_1} \sum_{k_3, k_4} g_{k_3 k_4} (\sigma_{m_3 m_4}^* + \pi_{m_3} \pi_{m_4}) + n \lambda_{j_1} \lambda_{j_2} \right] \end{aligned}$$

where  $\otimes$  is the Kronecker product, and as in (3.7)

$$B_1 = \sum_{i=1}^k t_i \text{diag}(c_{i1}, \dots, c_{ip})$$

$$G = (g_{ij})$$

$$\xi = (\mu_1, \mu_2, \dots, \mu_n)' \quad (6.5)$$

$$\pi = [I - \frac{2i}{\sqrt{n}} (I \otimes \Sigma)(G \otimes B_1)]^{-1} \xi = (\pi_1, \pi_2, \dots, \pi_{np})'$$

$$\Sigma^* = (I \otimes \Sigma) [I - \frac{2i}{\sqrt{n}} (G \otimes B_1)(I \otimes \Sigma)]^{-1} = (\sigma_{ij}^*)$$

and

$$i_1 = p \times (k_1 - 1) + j_1$$

$$m_1 = p \times (k_1 - 1) + j_1$$

$$i_2 = p \times (k_3 - 1) + j_1$$

$$m_2 = p \times (k_2 - 1) + j_1$$

$$i_3 = p \times (k_2 - 1) + j_2$$

$$m_3 = p \times (k_3 - 1) + j_2$$

$$i_4 = p \times (k_4 - 1) + j_2$$

$$m_4 = p \times (k_4 - 1) + j_2$$

Use the expansion that

$$[I - \frac{2i}{\sqrt{n}} (I \otimes \Sigma)(G \otimes B_1)]^{-1} = \sum_{s=0}^{\infty} [\frac{2i}{\sqrt{n}} (I \otimes \Sigma)(G \otimes B_1)]^s \quad (6.6)$$

$$[I - \frac{2i}{\sqrt{n}} (G \otimes B_1 \Sigma)]^{-\frac{1}{2}} = \exp \frac{1}{2} \left( \sum_{s=1}^{\infty} \frac{(2i)^s \text{tr } G^s \cdot \text{tr}(B_1 \Sigma)^s}{s \sqrt{n}^s} \right)$$

$$\text{and } \text{tr } \xi \xi' (G \otimes B_1) = \text{tr } B_1 U G U'.$$

We obtain the characteristic function as

$$\psi(t_1, \dots, t_k) = \exp(-\frac{1}{2} t' Q t)$$

$$\{1 + \frac{1}{\sqrt{n}} \sum_{i=1}^k i t_i (h_1 + h_2) + \frac{1}{\sqrt{n}} \sum_{i_1, i_2, i_3}^k (i^3 t_{i_1} t_{i_2} t_{i_3})\}$$

$$(h_3 + h_4 + h_5) + O(n^{-1}) \quad (6.7)$$

where

$$Q = (Q_{i_1 i_2}), \quad Q_{i_1 i_2} = 2 \frac{\text{tr} G^2}{n} \text{tr} R^{(i_1)} R^{(i_2)} + 4 \text{tr} R^{(i_1)} \psi^{(2, i_2)}$$

$$h_1 = \sum_{\alpha=1}^r \sum_{\beta \neq \alpha} \sum_{\substack{j_1 \in J_\alpha \\ j_2 \in J_\beta}} a_{i\alpha} \theta_{\alpha\beta}^{-1} \left( \frac{\text{tr} G^2}{n} (\sigma_{j_1 j_1} \sigma_{j_2 j_2} + \sigma_{j_1 j_2}^2) \right. \\ \left. + 2 \sigma_{j_1 j_2} T_{j_1 j_2}^{(2)} + \sigma_{j_1 j_1} T_{j_2 j_2}^{(2)} + \sigma_{j_2 j_2} T_{j_1 j_1}^{(2)} \right)$$

$$h_2 = \sum_{\alpha, \beta=1}^Y \sum_{\substack{j_1 \in J_\alpha \\ j_2 \in J_\beta}} a_{i\alpha\beta} \left( \frac{\text{tr} G^2}{n} \sigma_{j_1 j_2}^2 + 2 T_{j_1 j_2}^{(2)} \sigma_{j_1 j_2} \right)$$

$$h_3 = \frac{4}{3} \frac{\text{tr} G^3}{n} \text{tr} R^{(i_1)} R^{(i_2)} R^{(i_3)} + 4 \text{tr} R^{(i_1)} R^{(i_2)} \psi^{(3, i_3)} \quad (3.8)$$

$$h_4 = 4 \sum_{\alpha=1}^r \sum_{\beta \neq \alpha} \sum_{\substack{j_1 \in J_\alpha \\ j_2 \in J_\beta}} a_{i1\alpha} \theta_{\alpha\beta}^{-1} \left( \frac{\text{tr} G^2}{n} \varepsilon_{j_1 j_2}^{(i_2)} + \Omega_{j_1 j_2}^{(i_2)} + \Omega_{j_2 j_1}^{(i_2)} \right) \\ \left( \frac{\text{tr} G^2}{n} \varepsilon_{j_1 j_2}^{(i_3)} + \Omega_{j_1 j_2}^{(i_3)} + \Omega_{j_2 j_1}^{(i_3)} \right)$$

$$h_5 = 2 \sum_{\alpha, \beta} \sum_{\substack{j_1 \in J_\alpha \\ j_2 \in J_\beta}} a_{i\alpha\beta} \left( \frac{\text{tr} G^2}{n} \varepsilon_{j_1 j_1}^{(i_2)} + 2 \Omega_{j_1 j_1}^{(i_2)} \right) \\ \left( \frac{\text{tr} G^2}{n} \varepsilon_{j_2 j_2}^{(i_3)} + 2 \Omega_{j_2 j_2}^{(i_3)} \right)$$

$$T^{(2)} = \frac{u G^2 u'}{n}, \quad R^{(i)} = C^{(i)} \Sigma, \quad C^{(i)} = \text{diag}(c_{i1}, \dots, c_{ip})$$

$$\psi(j, i) = C^{(i)} \frac{u G^j u'}{n}, \quad \varepsilon^{(i)} = \Sigma R^{(i)}, \quad \Omega^{(i)} = T^{(2)} R^{(i)} \quad (6.9)$$

where  $A_{ij}$  denotes the  $(i, j)$ th elements of matrix  $A = (A_{ij})$ .

Inverting the characteristic function, we obtain the following expression for the asymptotic joint distribution of  $\underline{L} = (L_1, \dots, L_k)'$ .

$$\begin{aligned}
 f(L_1, \dots, L_k) &= N(\underline{L}, Q) \\
 &\times \left[ 1 + \frac{1}{\sqrt{n}} \sum_{i=1}^k H_i(\underline{L})(h_1 + h_2) \right. \\
 &\quad \left. + \frac{1}{\sqrt{n}} \sum_{i_1, i_2, i_3}^k H_{i_1 i_2 i_3}(\underline{L})(h_3 + h_4 + h_5) \right] + o(n^{-1})
 \end{aligned} \tag{6.10}$$

where  $N(\underline{L}, Q)$  and  $H_i(\underline{L})$ ,  $H_{i_1 i_2 i_3}(\underline{L})$  are as defined in Eq. (3.15)

## 7. Asymptotic Distributions of Functions of the Roots of the Complex Wishart Matrix

Let  $Z = Z_1 + i Z_2$  be a  $p \times n$  matrix and let the rows of  $(Z_1' : Z_2')$  be distributed independently as multivariate normal with covariance matrix

$$\begin{pmatrix} \Sigma_1 & \Sigma_2 \\ -\Sigma_2 & \Sigma_1 \end{pmatrix}$$

and let the mean vector of  $j$ -th row of  $(Z_1' : Z_2')$  be

$\underline{\mu}_j' = (\underline{\mu}_j^{(1)}, \underline{\mu}_j^{(2)})$ . Also let  $\tilde{S} = Z\bar{Z}'$  where  $\bar{Z}$  denotes the complex conjugate of  $Z$ . Then, the distribution of  $\tilde{S}$  is

known to be central or noncentral complex Wishart matrix

with  $\underline{n}$  degrees of freedom according as  $\tilde{M} = 0$  or  $\tilde{M} \neq 0$ , where  $\underline{M} = \underline{U}\underline{U}'$ ,  $E(Z) = \underline{U}$ . The expected value of  $S$  is given by

$$E(S) = 2n(\Sigma_1 - i\Sigma_2) + \tilde{M}$$

The matrix  $S$  is Hermitian and the eigenvalues of  $S/n$  are denoted by

$$\ell_1 \geq \dots \geq \ell_p.$$

In the sequel, we assume that  $E(\tilde{S}) = n \text{ diag. } (\lambda_1, \dots, \lambda_p)$  and  $\lambda_1 \geq \dots \geq \lambda_p$ . In addition, we assume that  $\lambda_i$ 's have multiplicity as in (3.1). Now, let

$$L_j = \sqrt{n} \{T_j(\ell_1, \dots, \ell_p) - T_j(\lambda_1, \dots, \lambda_p)\} \quad (7.1)$$

for  $j = 1, 2, \dots, k$  and the function  $T_j(\ell_1, \dots, \ell_p)$  satisfy the assumptions (3.2) and (3.3). Then, following the same lines

as in Section 3 for the real case, we obtain the following asymptotic expression for the joint density of  $L_1, L_2, \dots, L_k$ :

$$\begin{aligned} f_1(L_1, \dots, L_k) = & N(L, \tilde{Q}) \left[ 1 + \frac{1}{\sqrt{n}} \sum_{i=1}^k H_i(L) (\tilde{d}_1 + \tilde{d}_2) \right. \\ & + \frac{1}{\sqrt{n}} \sum_{i_1, i_2, i_3=1}^k H_{i_1 i_2 i_3}(L) (\tilde{g}_1 + \tilde{g}_2 + \tilde{g}_3) \left. \right] \\ & + O(n^{-1}) \end{aligned} \quad (7.2)$$

where

$$\begin{aligned} \tilde{Q} = (\tilde{Q}_{i_1 i_2}), \tilde{Q}_{i_1 i_2} = & 4 \operatorname{tr} \tilde{R}_1^{(i_1) \sim (i_2)} + 4 \operatorname{tr} \tilde{R}_2^{(i_1) \sim (i_2)} \\ & + 8 \operatorname{tr} \tilde{R}_1^{(i_1) \sim (i_2)}_{\psi} \end{aligned}$$

and  $\tilde{Q}$  is assumed to be nonsingular. Also,

$$\tilde{d}_1 = 4 \sum_{\alpha=1}^r \sum_{\beta \neq \alpha}^r \sum_{j_1 \in J_{\alpha}} \sum_{j_2 \in J_{\beta}} a_{i_{\alpha}} \theta_{\alpha \beta}^{-1} (\tilde{\sigma}_{j_1 j_1}^{(1)} \tilde{\sigma}_{j_2 j_2}^{(1)} + \tilde{\sigma}_{j_1 j_1}^{(1)} \tilde{v}_{j_2 j_2}^{(1)} + \tilde{\sigma}_{j_2 j_2}^{(1)} \tilde{v}_{j_1 j_1}^{(1)}) \quad (7.3)$$

$$\tilde{d}_2 = 2 \sum_{\alpha=1}^r \sum_{\beta=1}^r \sum_{j_1 \in J_{\alpha}} \sum_{j_2 \in J_{\beta}} a_{i_{\alpha \beta}} (\tilde{\sigma}_{j_1 j_2}^{(1)2} + 2 \tilde{\sigma}_{j_1 j_2}^{(1)} \tilde{v}_{j_1 j_2} - \tilde{\sigma}_{j_1 j_2}^{(2)2}) \quad (7.4)$$

$$\begin{aligned} \tilde{g}_1 = & \frac{8}{3} \operatorname{tr} \tilde{R}_1^{(i_1)} \tilde{R}_1^{(i_2)} \tilde{R}_1^{(i_3)} + 8 \operatorname{tr} \tilde{R}_1^{(i_1)} \tilde{R}_1^{(i_2)} \tilde{\psi}^{(i_3)} \\ & + 8 \operatorname{tr} \tilde{R}_2^{(i_1)} \tilde{R}_2^{(i_2)} \tilde{R}_1^{(i_3)} - 8 \operatorname{tr} \tilde{R}_2^{(i_1)} \tilde{R}_2^{(i_2)} \tilde{\psi}^{(i_3)} \end{aligned} \quad (7.5)$$

$$\tilde{g}_2 = 16 \sum_{\alpha=1}^r \sum_{\beta \neq \alpha}^r \sum_{j_1 \in J_{\alpha}} \sum_{j_2 \in J_{\beta}} a_{i_1 \alpha} \theta_{\alpha \beta}^{-1} [(\tilde{E}_{j_1 j_2}^{(i_2)} + \tilde{T}_{j_1 j_2}^{(i_2)} + \tilde{T}_{j_2 j_1}^{(i_2)} + \tilde{G}_{j_1 j_2}^{(i_2)})]$$

$$\begin{aligned}
& \times (\tilde{E}_{j_1 j_2}^{(i_3)} + \tilde{T}_{j_1 j_2}^{(i_3)} + \tilde{T}_{j_2 j_1}^{(i_3)} + \tilde{G}_{j_1 j_2}^{(i_3)} \\
& + (\tilde{U}_{j_1 j_2}^{(i_2)} - \tilde{U}_{j_2 j_1}^{(i_2)}) (\tilde{U}_{j_1 j_2}^{(i_3)} - \tilde{U}_{j_2 j_1}^{(i_3)}) \} \\
\end{aligned} \tag{7.6}$$

$$\begin{aligned}
\tilde{E}_3 = 8 \sum_{\alpha=1}^r \sum_{\beta=1}^r \sum_{j_1 \in J_\alpha} \sum_{j_2 \in J_\beta} a_{j_1 \alpha \beta} (\tilde{E}_{j_1 j_1}^{(i_2)} + 2\tilde{T}_{j_1 j_1}^{(i_2)} + \tilde{G}_{j_1 j_1}^{(i_2)}) \\
\times (\tilde{E}_{j_2 j_2}^{(i_3)} + 2\tilde{T}_{j_2 j_2}^{(i_3)} + \tilde{G}_{j_2 j_2}^{(i_3)})
\end{aligned} \tag{7.7}$$

where  $C^{(i)} = \text{diag}(c_{i1}, \dots, c_{ip})$ ,  $\Sigma_1 = (\tilde{\sigma}_{i_1 i_2}^{(1)})$ ,  $\Sigma_2 = (\tilde{\sigma}_{i_1 i_2}^{(2)})$

$$\tilde{M} = \sum_{j=1}^n (\mu_j^{(1)} \mu_j^{(1)'} + \mu_j^{(2)} \mu_j^{(2)'}) / 2n = (\tilde{v}_{j_1 j_2})$$

$$\tilde{R}_1^{(i)} = C^{(i)} \Sigma_1, \tilde{R}_2^{(i)} = C^{(i)} \Sigma_2, \tilde{\psi}^{(i)} = C^{(i)} \tilde{M}$$

$$\tilde{E}^{(i)} = \Sigma_1 C^{(i)} \Sigma_1, \tilde{G}^{(i)} = \Sigma_2 C^{(i)} \Sigma_2$$

$$\tilde{T}^{(i)} = \tilde{M} C^{(i)} \Sigma_1, \tilde{U}^{(i)} = \tilde{M} C^{(i)} \Sigma_2$$

Krishnaiah and Lee (1977) derived the asymptotic joint distributions of the linear combinations and ratios of the linear combinations of the eigenvalues of the central complex Wishart matrix when the roots are simple. These results are special cases of the results in this section.

## REFERENCES

- [1] Bryant, P. (1975). On testing for clusters using the sample covariance. J. Multivariate Anal. 5, 96-105.
- [2] Carter, E. M. and Srivastava, M. S. (1979). Asymptotic distributions of the latent roots of the non-central Wishart distribution and power of the L.R.T. for non-additivity. Technical Report No. 1, Department of Statistics, University of Toronto.
- [3] Corsten, L. C. A. and Van Eijnsbergen, A. C. (1972). Multiplicative effects in two-way analysis of variance. Statistica Neerlandica 26, 61-68
- [4] Fisher, R. A. (1939). The sampling distribution of some statistics obtained from non-linear equations. Ann. Eugenics 9, 238-249.
- [5] Fujikoshi, Y. (1978). Asymptotic expansions for the distributions of some functions of the latent roots of matrices in three situations. J. Multivariate Anal. 8, 63-72.
- [6] Ghosh, M. N. and Sharma, D. (1963). Power of Tukey's test for non-additivity. J. Roy. Statist. Soc. Ser. B 25, 213-219.
- [7] Gollob, H. F. (1968). A statistical model which combines features of factor analysis and analysis of variance techniques. Psychometrika 33, 73-116.
- [8] Johnson, D. E. and Graybill, F. A. (1972). An analysis of a two-way model with interaction and no replication. J. Amer. Statist. Assoc. 67, 862-868.
- [9] Kato, T. (1976). Perturbation Theory for Linear Operators. Springer-Verlag, New York.
- [10] Krishnaiah, P. R. and Chang, T. C. (1971). On the exact distributions of the extreme roots of the Wishart and MANOVA matrices. J. Multivariate Anal. 1, 108-117.
- [11] Krishnaiah, P. R. and Waikar, V. B. (1971). Simultaneous tests for equality of latent roots against certain alternatives - I. Ann. Inst. Statist. Math. 23, 451-468.
- [12] Krishnaiah, P. R. and Schuurmann, F. J. (1974). On the evaluation of some distributions that arise in simultaneous tests for the equality of the latent roots of the covariance matrix. J. Multivariate Anal. 4, 265-282.

- [13] Krishnaiah, P. R. and Lee, J. C. (1977). Inference on the eigenvalues of the covariance matrices of real and complex multivariate normal populations. In Multivariate Analysis-V (P. R. Krishnaiah, Ed.). North-Holland Publishing Company.
- [14] Krishnaiah, P. R. and Lee, J. C. (1979). On the asymptotic joint distribution of certain functions of the eigenvalues of four random matrices. J. Multivariate Anal. 9, 248-258.
- [15] Krishnaiah, P. R. (1980). Computations of some multivariate distributions. In Handbook of Statistics, Vol. 1 (P. R. Krishnaiah, Ed.). North-Holland Publishing Company.
- [16] Krishnaiah, P. R. and Yochmowitz, M. G. (1980). Inference on the structure of interactions in two-way classification model. In Handbook of Statistics, Vol. 1 (P. R. Krishnaiah, Ed.). North-Holland Publishing Company.
- [17] Lawley, D. N. (1956). Tests of significance for the latent roots of covariance and correlation. Biometrika 43, 128-136.
- [18] Mallows, C. L. (1961). Latent vectors of random symmetric matrices. Biometrika 48, 133-149.
- [19] Mandel, J. (1969). Partitioning the interaction in analysis of variance. Journal of Research of the National Bureau of Standards B 73, 3, 309-328.
- [20] Rao, C. R. (1965). Linear Statistical Inference and Its Applications. John Wiley & Sons.
- [21] Schuurmann, F. J., Krishnaiah, P. R. and Chattopadhyay, A. K. (1973). On the distributions of the ratios of the extreme roots to the trace of the Wishart matrix. J. Multivariate Anal. 3, 445-453.
- [22] Srivastava, M. S. and Carter, E. M. (1980). Asymptotic distribution of latent roots and applications. To appear in the Proceedings of the Halifax Conference on Multivariate Statistical Analysis.
- [23] Tukey, J. W. (1949). One degree of freedom for non-additivity. Biometrika 5, 232-242.
- [24] Williams, E. J. (1952). The interpretation of interactions in factorial experiments. Biometrika 39, 65-81.
- [25] Yochmowitz, M. G. and Cornell, R. G. (1978). Stepwise tests for multiplicative components of interaction. Technometrics 20, 79-84.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER <b>AFOSR-TR- 80 - 1164</b>	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Asymptotic Distributions of Functions of the Eigenvalues of the Real and Complex Noncentral Wishart Matrices		5. TYPE OF REPORT & PERIOD COVERED <i>Interim</i> Report
		6. PERFORMING ORG. REPORT NUMBER U.S. Stat. Rept. #80-11
7. AUTHOR(s) C. Fang and P. R. Krishnaiah		8. CONTRACT OR GRANT NUMBER(s) 49620-79-C-0161
9. PERFORMING ORGANIZATION NAME AND ADDRESS University of Pittsburgh Department of Mathematics & Statistics Pittsburgh, PA. 15260		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS <i>61102F 2304/AS</i>
11. CONTROLLING OFFICE NAME AND ADDRESS <i>AFOSR</i> <i>BOLLING AFB WASHINGTON DC 20332</i>		12. REPORT DATE July 1980
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES 38
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release: distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Asymptotic Distributions, Noncentral Wishart Matrix, Functions of Eigenvalues, Complex Matrices, Mixtures of Populations, Quadratic Forms, Perturbation Theory		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) In this paper, the authors derived asymptotic expressions for the joint densities of various functions of the noncentral real and complex Wishart matrices. Applications of the above results are also discussed in the area of reduction of dimensionality and testing for the structure of interaction in two way classification with one observation per cell.		

DD FORM 1 JAN 73 1473

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)